

Biomonitoring: Applications for Drinking Water Contaminants

Mark A. Werner, Ph.D.
Wisconsin Division of Public
Health

Presentation Goals

- Characterize the features and limitations of biomonitoring as an environmental public health tool
- Describe the use of biomonitoring to assess exposure in a population exposed to methyl-t-butyl ether (MtBE) in drinking water

Biomonitoring

- Assessment of human exposure to chemicals by measuring the chemicals or their metabolites in human specimens such as blood, urine, hair or nails
- Represents an increasingly available source of environmental public health data

How is Biomonitoring Useful?

- **Determine which chemicals are being absorbed and retained by humans**
- **Identify prevalence of people with levels above reference levels (Pb in blood)**
- **Establish reference ranges to identify what constitutes a significantly elevated level**

How is Biomonitoring Useful?

- **Assess effectiveness of efforts to reduce exposure to potentially harmful chemicals**
- **Determine distribution of exposure among at-risk populations (Hg among women of childbearing age)**
- **Identify trends in exposure over time**
- **Measure progress toward state and federal public health goals (HP 2010 and HW2010)**

Application to Drinking Water Contaminants

- Biomonitoring provides an *internalized* measure of exposure
 - For many contaminants, this provides a marker for exposure that logically would correlate better with actual exposure

Exposure to Drinking Water Contaminants

- Most commonly applied model:
 - $\text{Exposure (mg)} = \text{Sampled concentration (mg/L)} \times \text{Daily water intake (L)}$
 - Based on two critical assumptions:
 - Concentration at tap varies directly with sampled concentration
 - Absorbed dose varies directly with tap concentration

Tap Concentration vs. Sampled Concentration

- Why might these not correlate?
 - Effect of household plumbing (Pb, Cu)
 - Blending and distribution within a public water system
 - Treatment systems:
 - Water softeners (radium)
 - Whole-house treatment systems (iron, arsenic)
 - Reverse osmosis (most everything)

Absorbed Dose vs. Tap Concentration

- Why might these not correlate?
 - Use of other water sources
 - Variability in how much water people drink
 - Genetic differences in absorption & retention
 - Dietary, occupational and other exposures
 - Lead in children
 - Inhalation of chemicals
 - Radon in air vs. radon in water
 - Use of hot water and ventilation
 - Exposure to gasoline

Biomonitoring and Health Risk

- When are biomonitoring data a useful marker of health risk?
 - Substances with long biological half-lives
 - Lead, arsenic and cadmium
 - Conditions where exposure is great enough to allow for biomonitoring to exceed lab limits
 - Situations where chronic exposure is the concern
 - Probably not useful for nitrate

Biomonitoring Questions to Consider

- Can environmental exposure contribute to biological levels that can be measured in the medium of interest?
- Do biological levels better relate to health risk than sampled water concentrations?
- Does its use supplement information from water sampling in a meaningful way?

Biomonitoring Questions to Consider

- Does its use clarify who's at risk?
- Will people consent to sample extraction?
- Can the contribution of water-related exposure be quantified?

Human Exposure to Low Levels of MTBE in Drinking Water

Lisa Vallejo MPH

Lorraine Backer PhD MPH and Ben Blount PhD
National Center for Environmental Health
Centers for Disease Control and Prevention (CDC)

Mark Werner, PhD
Wisconsin Division of Public Health

Overview

- **MTBE from Public Health Perspective**
- **NHANES & National Report on Human Exposure to Environmental Chemicals**
- **Health Effects Studies**
- **Pilot Study**

Public Health Impact: MtBE Exposure

- **EPA has identified 106 cities/counties in non-attainment with the Clean Air Act**
 - 11 cities/counties in non-attainment for CO levels
- **USGS estimated 109 million Americans live in areas where MtBE is used**

NHANES

- **National Health and Nutrition Examination Survey**
 - Conducted by NCHS, CDC
- **Health and nutritional status**
- **Participants selected to represent the U.S. non-institutionalized, civilian population**

NHANES Subset 2000–2002: VOCs in Water and Blood

- **Subset of NHANES participants**
 - Random selection of >1000 NHANES participants
 - 1025 blood and 1256 water samples
 - Analysis of VOCs, including MtBE

NHANES Subset 2000-2002: MtBE Results

	Blood	Water
Geometric Mean	74 PPTr	190 PPTr
Median	14 PPTr	80 PPTr
Sample Size	1025	1256


MtBE and Health Effects

- **Acute health effects**
- **Chronic health effects**
- **Animal study**



Acute Health Effects


Study A



Source: Moolenaar et al., 1994. Archives of Environmental Health.

Acute Health Effects

Study A



Source: Moolenaar et al., 1994. Archives of Environmental Health.

Acute Health Effects

Study B

- Conducted in Stamford, Connecticut during oxygenated fuel program in 1993
 - Measured MtBE in personal breathing zone air, blood
 - Symptom questionnaire
- Results
 - MtBE levels in blood correlated with PBZ air
 - Symptom reporting associated with higher levels of MtBE in blood (2.4 ug/L)

Source: White et al., 1995. Archives of Environmental Health.

Chronic Health Effects

- 60 individuals exposed to 76 PPB to 14 PPB of MtBE & benzene in drinking water for several years
- Reported apoptosis in leukocytes in individuals exposed to MtBE and benzene in drinking water
- Limitation: Simultaneous exposure to benzene

Source: Vojdani et al., 1997. Human & Experimental Toxicology.

Animal Study

- Sprague-Dawley rat studies of MtBE
 - Doses: 1000; 250 ; & 0 mg/kg body weight
- Increase in leydig cell tumors in male rats
- Increase in lymphomas and leukemias in female rats

Source: Belpoggi et al., 1995. Toxicology & Industrial Health.

MtBE Exposure

What We Know

- **Potential widespread exposure from gasoline emissions and drinking water**
 - Possible carcinogen
 - Possible apoptosis-inducer



What's the Public Health Risk?

- **Verify extent of exposure**
- **Verify published results in a larger human study**

Objective of Pilot Study

To determine whether we could detect measurable levels in the blood of people exposed to low levels of MTBE in their drinking water


Pilot Study

- **Small community in Wisconsin**
- **Water contaminated with 1.2 - 5.0 ppb MtBE**
- **9 adult participants**
- **Collected blood and water samples**
 - November 2002 (Well with no MtBE)
 - December 2002 (Well with MtBE)

Study Inclusion Criteria

- Adults
- Drink at least two 8-oz glasses of tap water each day
- Could not pump gas or visit gasoline station 48 hours prior to blood draw
- Consent to blood draw and exposure questionnaire


Methods




Sample Collection

- **Blood Samples**
 - Vacutainer® tubes prepared by CDC
 - Certified phlebotomist
 - 10-mL blood sample
 - Shipped overnight to CDC
- **Water Samples**
 - Non-aerated household faucets
 - 12-mL glass vials
 - Faucet ran for 3 minutes
 - Vial slightly overfilled
 - Shipped overnight to CDC


Methods



Headspace SPME




Methods




Desorbing the SPME Fiber

Analytes are baked off (desorbed) the SPME fiber and swept onto the GC column.




Methods




Results Demographics

- **Average Age = 43 years**
- **All White, Non-Hispanic**
- **4 Females and 2 Males**


Results Blood



Results Blood




Results Water



Results


Water



Discussion

- **MTBE metabolism**
 - Oral exposure
- **Sample population**
 - Sample size
 - Minimal commuting

MTBE in Blood Following Oral Exposure 250 mL of 13 ppm



Conclusions

- **Very low levels of MTBE were detected in the blood of people exposed to tap water contaminated with very low levels of MTBE**
- **Blood levels of MTBE can be used as a biological marker of exposure in drinking water**

Next Steps for MTBE

- **Conduct epidemiologic study of health effects from exposure to MTBE**
- **Assess how different exposure routes impact internal dose and health effects**

Acknowledgements

CDC: Lisa Vallejo, Lorrie Backer, Ben Blount, David Ashley, Fred Cardinali, Lalith Silva
Sauk County Health Department